

Ordering ambiguity versus representation

A. de Souza Dutra^{a,b*}

^aAbdus Salam ICTP, Strada Costiera 11, 34014 Trieste Italy

^bUNESP-Campus de Guaratinguetá-DFQ[†]

Av. Dr. Ariberto Pereira da Cunha, 333

C.P. 205

12516-410 Guaratinguetá SP Brasil

February 1, 2008

Abstract

In this work we show that the ordering ambiguity on quantization depends on the representation choice. This property is then used to solve unambiguously a particular system. Finally we speculate on the consequences for more involved cases.

PACS numbers: 03.65.Ca

*E-mail: dutra@feg.unesp.br

[†]Permanent Institution

The problem of ordering ambiguity is one of the long standing questions of the quantum mechanics. This question has attracted the attention of some of the founders of the quantum mechanics. Born and Jordan, Weyl, Dirac and von Newmann worked on this matter, as can be verified from the excellent review by Shewell [1]. This is a hard problem which has advanced very few along the last decades. Notwithstanding, as a consequence of its importance in some experimental situations like impurities in crystals [2]-[4], the dependence of nuclear forces on the relative velocity of the two nucleons [5], [6], and more recently the study of semiconductor heterostructures [7],[8], the interest in such kind of systems never vanished. Furthermore, taking in account the spatial variation of the semiconductor type, some effective Hamiltonians were proposed with a spatially dependent mass for the carrier [9]-[14]. Some time ago we discussed the exact solvability of some classes of Hamiltonians with ordering ambiguity [15]. In fact, the problem of the spatially dependent mass is presenting a growing interest along the last few years [15]-[30].

Let us present the idea we are interested in develop in this work by illustrating it through the study a concrete example. In [15] it was shown that for a system whose quantum Hamiltonian has as one of its parts an operator version of the classical function $f(x)$ p . In the coordinate representation its operator counterpart can be written as

$$f(x) p \rightarrow \frac{f^\alpha(\hat{x}) \hat{p} f^\beta(\hat{x}) + f^\beta(\hat{x}) \hat{p} f^\alpha(\hat{x})}{2}, \quad (1)$$

where $\alpha + \beta \equiv 1$. By using the usual coordinate representation for the operator $\hat{p}$, and manipulating the above operator in order to put it at right, one can see that one obtains for instance

$$f^\alpha(\hat{x}) \hat{p} f^\beta(\hat{x}) = f(\hat{x}) \hat{p} - i \hbar \beta \frac{df(\hat{x})}{d\hat{x}}. \quad (2)$$

Now, using the corresponding operator for $f^\beta(\hat{x}) \hat{p} f^\alpha(\hat{x})$, and then calculating the Hermitian operator (1) with these features, one gets finally

$$\frac{f^\alpha(\hat{x}) \hat{p} f^\beta(\hat{x}) + f^\beta(\hat{x}) \hat{p} f^\alpha(\hat{x})}{2} = f(\hat{x}) \hat{p} - \frac{i \hbar}{2} \frac{df(\hat{x})}{d\hat{x}}, \quad (3)$$

from which we conclude that there is no ordering ambiguity in this representation and any ordering used will conduce essentially to the same final

answer, as observed in [15]. However, despite of being an important case of ordering, due to its application in the case of the minimal gauge coupling, it can not be used itself as a Hamiltonian, at least an usual one, because the momentum appears linearly in it. At this point we introduce the main idea underlying this work, by remembering that one could interchange the role of x and p , and discussing the case of the quantization of the classical function $g(p)$ in the momentum representation. It is not hard to conclude, through an absolutely analogous analysis in the momentum representation that the Hermitian quantization of this function is unambiguous, and looks like

$$\frac{g^\alpha(\hat{p}) \hat{x} f^\beta(\hat{p}) + f^\beta(\hat{p}) \hat{x} f^\alpha(\hat{p})}{2} = f(\hat{p}) \hat{x} + \frac{i\hbar}{2} \frac{df(\hat{p})}{d\hat{p}}. \quad (4)$$

Note, however, that this operator is surely ambiguous in the coordinate representation. From the above calculation we can conclude that the ordering ambiguity has a dependence on the choice of representation and, as far we know, this feature was not taken into account in the literature up to now. Furthermore, this last operator can be thought as a Hamiltonian if we choose $g(\hat{p}) = \hat{p}^2$. In this special case, we would have a system with spatial dependence in the spatial coordinate ($m(x) \sim \frac{1}{x}$). This is an example of Hamiltonian which is ambiguous in the coordinate representation and not in the momentum one. In cases like this, one could calculate the wave function in the momentum representation and then transform it through

$$\psi(x, t) = \int dp \tilde{\psi}(p, t) e^{\frac{i}{\hbar} p x}, \quad (5)$$

to the coordinate representation if necessary.

For the sake of concreteness, from now on we discuss this case with more details. Firstly, the time-independent wave function equation in the momentum representation is given by

$$i\hbar p^2 \frac{d\tilde{\psi}(p)}{dp} + i\hbar p \tilde{\psi}(p) = E \tilde{\psi}(p). \quad (6)$$

After a straightforward calculation, one obtains for the unambiguous wave function in the momentum representation

$$\tilde{\psi}(p) = N \frac{e^{\left(\frac{i}{\hbar} \frac{E}{p}\right)}}{p}, \quad (7)$$

where N is an arbitrary integration constant. We can now calculate its Fourier transform, in order to obtain the corresponding coordinate representation wave function. So, we must perform the following integration

$$\psi(x) = N \int_{-\infty}^{\infty} \frac{dp}{p} e^{\left(\frac{i}{\hbar} \frac{E}{p} + px\right)}. \quad (8)$$

In order to reach this goal, we separate the integral in two sectors, that for positive p and that for negative ones. So we gets

$$\psi(x) = N \left\{ \int_{-\infty}^0 \frac{dp}{p} e^{-\left(\frac{i}{\hbar} \frac{E}{p} + px\right)} + \int_0^{\infty} \frac{dp}{p} e^{\left(\frac{i}{\hbar} \frac{E}{p} + px\right)} \right\}, \quad (9)$$

which after some manipulations can be rewritten as

$$\psi(x) = 2iN \int_0^{\infty} \frac{dp}{p} \sin\left(\frac{i}{\hbar} \frac{E}{p} + px\right). \quad (10)$$

Then, after using usual trigonometric identities and the known result

$$\int_0^{\infty} \frac{du}{u} \sin(au) \cos\left(\frac{b}{u}\right) = \frac{\pi}{2} J_0\left(2(a^2 b^2)^{\frac{1}{4}}\right) = \int_0^{\infty} \frac{du}{u} \sin\left(\frac{b}{u}\right) \cos(au), \quad (11)$$

where $J_0(z)$ is the Bessel function of first kind. One obtains finally that

$$\psi(x) = 2\pi i N J_0\left(\frac{2}{\hbar} \sqrt{|Ex|}\right). \quad (12)$$

Have we started in the coordinate representation, the wave function equation to be solved would be

$$x^2 \frac{d^2 \psi(x)}{dx^2} + x \frac{d\psi(x)}{dx} - \alpha \gamma \psi(x) = -\left(\frac{E}{\hbar^2}\right) x \psi(x), \quad (13)$$

where we ordered the operator coming from $x p^2$ using

$$O_p \equiv \frac{1}{2} \left(\hat{x}^\alpha \hat{p} \hat{x}^\beta \hat{p} \hat{x}^\gamma + \hat{x}^\gamma \hat{p} \hat{x}^\beta \hat{p} \hat{x}^\alpha \right) = \hat{x} \hat{p}^2 - i\hbar \hat{p} + \alpha \gamma \hat{x}^{-1}, \quad (14)$$

and we used $\alpha + \beta + \gamma = 1$. It can be noted that, if make the variable transformation $|x| = \frac{\hbar^2}{4E} w$, the above equation can be cast in the form

$$w^2 \frac{d^2 \psi}{dw^2} + w \frac{d\psi}{dw} + (w^w - 4\alpha\gamma) \psi = 0, \quad (15)$$

which is the differential equation of the first kind Bessel function. So, we get finally that, in the coordinate representation the ambiguous wave function is expressed as

$$\psi(x) = N J_{\alpha\gamma} \left(\frac{2}{\hbar} \sqrt{|E x|} \right), \quad (16)$$

once E is positive definite. We conclude that the compatibility of the solutions coming from the two representations, imposes us to fix one of the parameters appearing in the index of the Bessel function ($\alpha = 0$ or $\gamma = 0$) in the coordinate representation. As a consequence of the symmetry between these parameters in the operator definition, is equivalent to choose any of them equal to zero. Choosing to make $\gamma = 0$, we conclude that $\beta = 1 - \gamma$, and we ends with a subclass of operators, compatibles in both representations,

$$O_\alpha = \frac{1}{2} \left(\hat{x}^\alpha \hat{p} \hat{x}^{1-\alpha} \hat{p} + \hat{p} \hat{x}^{1-\alpha} \hat{p} \hat{x}^\alpha \right). \quad (17)$$

Note that the case of the Li and Khun ordering [13], which we have shown to be equivalent to the Weyl ordering [15], corresponds to the choice $\alpha = \frac{1}{2}$.

Below we are going to prove that, in fact, there is no remaining ambiguity because all choices of α are equivalent. For this we note that

$$\hat{x}^\alpha \hat{p} \hat{x}^{1-\alpha} \hat{p} = \sqrt{\hat{x}} \hat{p} \sqrt{\hat{x}} \hat{p} + i \hbar \left(\alpha - \frac{1}{2} \right) \hat{p}; \quad (18)$$

$$\hat{p} \hat{x}^{1-\alpha} \hat{p} \hat{x}^\alpha = \hat{p} \sqrt{\hat{x}} \hat{p} \sqrt{\hat{x}} - i \hbar \left(\alpha - \frac{1}{2} \right) \hat{p},$$

so that the operator O_α is simply rewritten as

$$O_\alpha = \frac{1}{2} \left(\sqrt{\hat{x}} \hat{p} \sqrt{\hat{x}} \hat{p} + \hat{p} \sqrt{\hat{x}} \hat{p} \sqrt{\hat{x}} \right) = O_{Weyl}. \quad (19)$$

So we have finally demonstrated that, at least for this particular case, we have been able to avoid the ordering ambiguity by working in the momentum representation. In fact, this conclusion is still true if we include a binding potential energy in the original Hamiltonian. One can show also that, for a given class of potentials, the problem can be even exactly solvable. Furthermore we have shown that this unambiguous quantization corresponds to the so called Weyl ordering.

It is interesting to see that, in some very recent papers, it was adjudicate in favor of a Schroedinger equation in a phase-space representation, where

appears a very interesting kind of mixing between the usual coordinate and momentum representations [31],[32]. It would be very interesting to see if this generalized representation could be useful in some particular problem, where there exist the with usual ordering ambiguity both in coordinate and momentum representations and, maybe not in this new representation.

Acknowledgments: The author is grateful to CNPq for partial financial support and to the professor M. Hott for many helpful discussions. This work has been done during a visit under the auspices of the Associate Scheme of the Abdus Salam ICTP.

References

- [1] J. R. Shewell, Am. J. Phys. **27** (1959) 16.
- [2] J. M. Luttinger and W. Kohn, Phys. Rev. **97** (1955) 869.
- [3] G. H. Wannier, Phys. Rev. **52** (1957) 191.
- [4] J. C. Slater, Phys. Rev. **76** (1949) 1592.
- [5] Ó. Rojo and J. S. Levinger, Phys. Rev. **123** (1961) 2177.
- [6] M. Razavy, G. Field and J. S. Levinger, Phys. Rev. **125** (1962) 269.
- [7] G. Bastard, Wave Mechanics Applied to Semiconductor Heterostructures, Les Éditions de Physique, Les Ullis, 1992.
- [8] C. Weisbuch and B. Vinter, Quantum Semiconductor Heterostructures, Academic Press, New York, 1993.
- [9] D. J. BenDaniel and C. B. Duke, Phys. Rev. B **152** (1966) 683.
- [10] Q. G. Zhu and H. Kroemer, Phys. Rev. B **27** (1983) 3519.
- [11] T. Gora and F. Williams, Phys. Rev. **177** (1969) 1179.
- [12] G. Bastard. Phys. Rev. B **24** (1981) 5693.
- [13] T. L. Li and K. J. Kuhn, Phys. Rev. B **47** (1993) 12760.
- [14] F. S. A. Cavalcante, R. N. Costa Filho, J. Ribeiro Filho, C. A. S. de Almeida and V. N. Freire, Phys. Rev. B **55** (1997) 1326.
- [15] A. de Souza Dutra and C. A. S. de Almeida, Phys. Lett. A **275** (2000) 25.
- [16] A. de Souza Dutra, M. B. Hott and C. A. S. Almeida, Europhys. Lett. **62** (2003) 8.
- [17] F. S. A. Cavalcante, R. N. Costa Filho, J. Ribeiro Filho, C. A. S. de Almeida and V. N. Freire, Phys. Rev. B **55** (1997) 1326.
- [18] G. Chen, Chin. Phys. **14** (2005) 460.

- [19] G. Chen and Z. D. Chen, Phys. Lett. A **331** (2004) 312.
- [20] A. A. Stahlhofen, J. Phys. A **37** (2004) 10129-10138,
- [21] B. Bagchi, P. Gorain, C. Quesne and R. Roychoudhury, Mod. Phys. Lett. A **19** (2004) 2765-2775,
- [22] K. Bencheikh, K. Berkane and S. Bouizane, J. Phys. A **37** (2004) 10719-10725,
- [23] J. A. Yu, S. H. Dong, Phys. Lett. A **325** (2004) 194-198,
- [24] C. Quesne and V. M. Tkachuk, J. Phys. A **37** (2004) 4267-4281,
- [25] Y. C. Ou, Z. Q. Cao and Q. H. Shen, J. Phys. A **37** (2004) 4283-4288,
- [26] R. Koc and H. Tutunculer, Annalen der Physik **12** (2003) 684-691,
- [27] A. D. Alhaidari, Phys. Rev. A **66** (2002) 042116,
- [28] B. Roy and P. Roy, J. Phys. A **35** (2002) 3961-3969.
- [29] S. Ramgoolam, B. Spence and S. Thomas, Nucl. Phys. B **703** (2005) 236.
- [30] A. de Souza Dutra, M. B. Hott and V. G. C. S. dos Santos, Europhys. Lett. **71** (2005) 166.
- [31] Q. S. Li, G. M. Wei and L. Q. Lü, Phys. Rev A **70** (2004) 022105.
- [32] D. Chruściński and K. Młodawski, Phys. Rev. A **71** (2005) 052104.